

Semiclassical Theory of Chaotic Conductors

Stefan Heusler¹, Sebastian Müller¹, Petr Braun^{1,2}, Fritz Haake¹

¹*Fachbereich Physik, Universität Duisburg-Essen, 45117 Essen, Germany*

²*Institute of Physics, Saint-Petersburg University, 198504 Saint-Petersburg, Russia*

We calculate the Landauer conductance through chaotic ballistic devices in the semiclassical limit, to all orders in the inverse number of scattering channels without and with a magnetic field. Families of pairs of entrance-to-exit trajectories contribute, similarly to the pairs of periodic orbits making up the small-time expansion of the spectral form factor of chaotic dynamics. As a clue to the exact result we find that close self-encounters slightly hinder the escape of trajectories into leads.

PACS numbers: 73.23.-b, 05.45.Mt, 03.65.Sq, 73.20.Fz, 72.20.My

Electron transport through conductors is usually dominated by scattering from disordered impurities. The pertinent theory involves averages over random configurations of scatterers and ultimately amounts to random-matrix theory. Typical consequences of disorder on transport, like weak localization, actually survive for “clean” systems whose linear dimension is smaller than the mean free path, provided the ballistic motion is chaotic due to irregularly shaped boundaries of the “cavity” serving as the conductor [1, 2]. The equivalence of chaos in the ballistic limit and impurity scattering represents a theoretical challenge similar to that of proving the so-called Bohigas-Giannoni-Schmit conjecture about universality of fluctuations in energy spectra of chaotic dynamics.

Gutzwiller’s semiclassical periodic-orbit theory has recently been brought to success in explaining spectral fluctuations of individual chaotic dynamics as faithful to the ensemble averages provided by random-matrix theory [3]. Analogous attempts have been undertaken for ballistic transport [2, 4], employing semiclassical techniques in Landauer’s scattering approach to conductance [5]. The two lowest-order terms in a $1/N$ -expansion (with N the dimension of the scattering matrix) of the conductance were thus found in agreement with RMT. Here we take up that line of thought, with two crucial modifications to be explained below, which yield the conductance to all orders in $1/N$. Both for systems with and without time reversal invariance as well as for the crossover between these two cases caused by a magnetic field we confirm the equivalence to impurity scattering as predicted by random-matrix theory. Each order of the $1/N$ expansion turns out one-to-one with the small-time expansion of the universal spectral form factor given in [3]. Our confirmation of the validity of the RMT conductance for individual chaotic cavities with arbitrary N is particularly relevant for (electronic or microwave) experiments involving a number of channels of order unity [6].

We imagine a two dimensional cavity connected to the outside by two leads respectively supporting N_1 and N_2 channels. The entrance-to-exit transition amplitudes for (electron) waves then form a unitary $N \times N$ S -matrix where $N = N_1 + N_2$. Sub-blocks t, t' of respective size $N_1 \times N_2$, $N_2 \times N_1$ comprise the amplitudes for transitions between different leads while the remaining sub-blocks

r, r' refer to “reflections”, i. e. transitions with entrance and exit at the same lead. We are concerned with the conductance between different leads, given by Landauer [5] as $G = \text{Tr}(t^\dagger t)$, in units of the quantum conductance $G_0 = e^2/\pi\hbar$. Random-matrix theory yields an averaged conductance G_β , the average being over the Wigner-Dyson circular ensembles of unitary matrices from the orthogonal ($\beta = 1$) or unitary ($\beta = 2$) class [7, 8],

$$G_\beta = \frac{N_1 N_2}{N - 1 + 2/\beta} = \frac{N_1 N_2}{N} \left\{ 1 + \sum_{n=1}^{\infty} \frac{(1 - 2/\beta)^n}{N^n} \right\}. \quad (1)$$

We first deal with the orthogonal and unitary classes. Further below, we shall allow for a magnetic field causing a smooth transition between these classes. Random-matrix theory provides an integral expression [9] $G(\xi)$ interpolating between G_1 and G_2 , as a parameter ξ proportional to the squared magnetic field is varied.

We shall derive a semiclassical expression for the conductance of a single chaotic cavity. The van Vleck propagator for the amplitudes t brings classical entrance-to-exit trajectories into play [1, 2], $t_{nm} \propto \sum_\gamma A_\gamma e^{iS_\gamma/\hbar}$, with S_γ the action and A_γ a stability amplitude of γ ; the sum includes all trajectories connecting the leads with discrete incident angles, one for each channel. The Landauer conductance becomes the double sum [2],

$$\text{Tr}(t^\dagger t) = \frac{1}{T_H} \left\langle \sum_{\gamma, \gamma'} A_\gamma A_{\gamma'}^* e^{i(S_\gamma - S_{\gamma'})/\hbar} \right\rangle, \quad (2)$$

somewhat similar to Gutzwiller’s [3] double sum over periodic orbits for the spectral form factor; here T_H is the Heisenberg time, related to the volume of the energy shell as $T_H = \Omega/2\pi\hbar$. The brackets $\langle \cdot \rangle$ denote an energy average over a small interval near the Fermi energy.

In the limit $\hbar \rightarrow 0$ only trajectory pairs with the action difference $\Delta S = S_\gamma - S_{\gamma'}$ of the order $\hbar$ may survive the averaging implied in (2). The simplest such family is the diagonal one with $\gamma = \gamma'$. To evaluate the diagonal approximation $\text{Tr}(t^\dagger t)_{\text{diag}} = T_H^{-1} \langle \sum_{\gamma=\gamma'} |A_\gamma|^2 \rangle$, Richter and Sieber [4] invoked a new sum rule which we write as

$$\text{Tr}(t^\dagger t)_{\text{diag}} = \frac{1}{T_H} \left\langle \sum_{\gamma=\gamma'} |A_\gamma|^2 \right\rangle = \frac{N_1 N_2}{T_H} \int_0^\infty dT e^{-\frac{T}{T_H}} T$$




FIG. 1: Pairs of trajectories γ, γ' contributing to the conductance: a) a Richter/Sieber pair differing in a 2-encounter, b) a pair of trajectories differing in one 2-encounter and one 3-encounter. In a), γ and γ' differ in the sense of traversal of one loop ($\nu = 1$), and the relative orientations of stretches in the 2-encounter lead to $\mu = 0$. For b) we have $\nu = 3$; $\mu = -1$ for the 2-encounter and $\mu = 1$ for the 3-encounter

$$= \frac{N_1 N_2}{N}. \quad (3)$$

Like the somewhat analogous sum rule given by Hanay and Ozorio de Almeida [10] for periodic orbits, the Richter-Sieber sum reflects the ergodicity of long trajectories; in addition, it accounts for the escape of a trajectory hitting a lead after a dwell time T by the exponential “survival probability” $e^{-\frac{N}{T_H}T}$. In spite of its quantum guise, which is convenient for our purposes, the escape rate N/T_H is, of course, a classical quantity; Planck’s constant cancels since the number of channels is proportional to the ratio of the width of the leads and the de Broglie wavelength. Clearly, the diagonal approximation gives the first term of the $1/N$ expansion of the random-matrix conductance (1), and even the full conductance for the unitary symmetry class.

Higher-order terms are due to families of pairs (γ, γ') of trajectories. The partner trajectories in a pair nearly coincide, at least up to time reversal, in “loops” whose cumulative duration is of the order T_H ; these loops are separated by close self-encounters of each trajectory; the encounter stretches have durations of the order of the Ehrenfest time $T_E \propto \ln \hbar \ll T_H \propto 1/\hbar$ and connect the loops differently for γ and γ' . The various families of pairs (γ, γ') can be imagined as arising from the pairs of periodic orbits contributing to the analogous expansion of the spectral form factor [3] by cutting a loop into two and pulling the two open ends to different leads. The cutting described actually turns the families of pairs of periodic orbits giving the term τ^n in the τ expansion of the spectral form factor into the pairs of trajectories contributing to the n -th order in the $1/N$ expansion of the conductance. The dwell time T of γ and γ' is then the cumulative duration of all loops and encounter stretches.

A clue to the full expansion in search is the survival probability. That quantity is a little larger than $e^{-\frac{N}{T_H}T}$ for all trajectory pairs except the “trivial” ones, $\gamma' = \gamma$. To see that, we focus on, say, an l -encounter where l

stretches of γ (and of γ') are mutually close at least up to time reversal and separated from the leads by loops; each of the l intra-encounter stretches has the duration t_{enc} . Survival of γ inside the cavity during the first of the l stretches implies survival during the $l - 1$ other stretches. The whole encounter therefore does not contribute lt_{enc} but only t_{enc} to the effective exposure time in the exponential survival probability. The cumulative exposure time is the sum of all loop durations, augmented by one instead of l contributions t_{enc} for each l -encounter, $T_{\text{exp}} = \sum_{\text{loops}} t_{\text{loop}} + \sum_{\alpha} t_{\text{enc}}^{\alpha} = T - \sum_{\alpha} (l_{\alpha} - 1)t_{\text{enc}}^{\alpha}$, the index α labeling the encounters. Due to $T_E \ll T_H$ the survival probability $e^{-\frac{N}{T_H}T_{\text{exp}}}$ differs only little from the naively expected value $e^{-\frac{N}{T_H}T}$, but the difference will presently be seen to be decisively important.

We illustrate our procedure for the first correction to the diagonal approximation which involves partner trajectories differing in a single 2-encounter. Fig. 1a depicts the two encounter stretches and three loops with durations $t_i, i = 1, 2, 3$; the dwell time is $T = t_1 + t_2 + t_3 + 2t_{\text{enc}}$. The two encounter stretches must be nearly mutually time reversed since otherwise the partner trajectory would decompose such that one of the loops would connect the two leads while the other would close in on itself to form a periodic orbit; this is why the first off-diagonal correction requires time reversal invariance.

We may search for 2-encounters drawing a Poincaré section $\mathcal{P}$ transverse to γ . If the section hits a 2-encounter, γ must pierce through $\mathcal{P}$ in two points which are almost mutually time reversed. We denote the stable and unstable shifts of the second (time reversed) piercing relative to the first by s, u . For the quantitative definition of 2-encounters we require $|u| \leq c, |s| \leq c$ with c a constant independent of $\hbar$ (whose exact value is irrelevant). The coordinates u, s determine the encounter duration $t_{\text{enc}} = (1/\lambda) \ln c^2/|su|$ and the action difference $\Delta S = S_{\gamma} - S_{\gamma'} = us$; here λ is the Lyapunov constant.

The ergodic probability of the second piercing occurring with given s, u and duration t_2 of the loop

separating the encounter stretches is $ds du dt_2/\Omega$. We can write the number of trajectory pairs generated by 2-encounters with the given action difference ΔS as $d\Delta S \int_{|s|<c, |u|<c} ds du w_T(u, s) \delta(\Delta S - us)$. The weight function $w_T(u, s)$ is obtained from the ergodic return density $1/\Omega$ by (i) integrating over all durations t_2 of the loop separating the encounter stretches; (ii) integrating over all positions of the Poincaré section which is equivalent to integrating over the duration of the first loop t_1 ; (iii) dividing by the encounter duration t_{enc} (the probability to cross an encounter is proportional to its duration; in order to obtain the number of encounters from the crossing probability the encounter duration has to be divided out). Requiring all loop times to be positive we get

$$\begin{aligned} w_T(u, s) &= \int \frac{dt_1 dt_2}{\Omega t_{\text{enc}}} \Theta(t_1) \Theta(t_2) \Theta(T - t_1 - t_2 - 2t_{\text{enc}}) \\ &= \frac{(T - 2t_{\text{enc}}(u, s))^2}{2t_{\text{enc}}(u, s)\Omega}. \end{aligned} \quad (4)$$

The first off-diagonal term in the conductance thus reads

$$G^{(1)} = \left\langle \int_{|s|<c, |u|<c} ds du e^{isu/\hbar} \sum_{\gamma} \frac{|A_{\gamma}|^2}{T_H} w_T(s, u) \right\rangle. \quad (5)$$

The sum over γ is now calculated using the rule (3) with the crucial modification pointed out above, i.e. the survival probability $e^{-\frac{N}{T_H}(t_1+t_2+t_3+t_{\text{enc}})}$. The integrations over t_1, t_2 (in w_T) and over the dwell time T can be replaced by integrations over all three loop durations. The lower limit of a loop duration is *zero* which expresses a second crucial ingredient of our approach: *the encounter stretches may neither overlap nor stick out of the cavity openings* [11]; the upper limit is infinity. The result

$$\begin{aligned} G^{(1)} &= \frac{N_1 N_2}{T_H} \left(\prod_{i=1}^3 \int_0^\infty dt_i e^{-\frac{N}{T_H} t_i} \right) \\ &\times \left\langle \int_{|s|<c, |u|<c} ds du \frac{e^{-\frac{N}{T_H} t_{\text{enc}} + isu/\hbar}}{\Omega t_{\text{enc}}} \right\rangle \rightarrow -\frac{N_1 N_2}{N^2} \end{aligned} \quad (6)$$

obtains a factor T_H/N from each loop-time integration. To calculate the integral over s, u we expanded in powers of the encounter duration, $e^{-\frac{N}{T_H} t_{\text{enc}}} = 1 - \frac{N}{T_H} t_{\text{enc}} + \dots$; the contribution of the zero-order term is annihilated by averaging; terms of second and higher order vanish in the limit $\hbar \rightarrow 0$ [3]; in the remaining first-order term t_{enc} cancels against the denominator whereupon the integral becomes trivial, $\frac{1}{\Omega} \int_{-c}^c ds du e^{isu/\hbar} \rightarrow 2\pi\hbar/\Omega = 1/T_H$.

Higher orders in $1/N$ are due to pairs (γ, γ') generated by reshuffling intra-encounter connections of loops within an l -encounter with $l > 2$ or within several encounters; see Fig. 1b. In general, such reconnections arise in v_l l -encounters, $l = 2, 3, \dots$; we collect these multiplicities in a vector $\vec{v}$. The total number of encounters is $V = \sum_{l=2}^\infty v_l$ while $L = \sum_{l=2}^\infty l v_l$ gives the number of the encounter stretches; the total number of loops is $L + 1$.

Each encounter requires its own Poincaré section which latter contains l close-by piercings by γ ; the separations of subsequent piercings with respect to the first ones can be expressed in terms of coordinates $s_j, u_j, j = 1, \dots, l-1$ obeying $|s_j|, |u_j| \leq c$ [3]. The ergodic probability density for the $L - V$ returns is $1/\Omega^{L-V}$. To obtain the weight function analogous to (4) for the number of encounter sets with a given $\vec{v}$, the ergodic density has to be integrated over all loop times but one, and divided by the product of all encounter durations. In addition the result has to be multiplied by the number $\mathcal{N}(\vec{v})$ of topological versions (“structures”) of trajectory pairs differing by (i) the order in which encounters are visited in γ ; (ii) the relative direction of stretches in each encounter and (iii) the intra-encounter reconnections leading from γ to γ' . The action difference in the pair can be written as $\Delta S = \sum_{\alpha, j} s_{\alpha j} u_{\alpha j}$. When applying the sum rule we use the exposure time $T_{\text{exp}} = \sum_{i=1}^{L+1} t_i + \sum_{\alpha=1}^V t_{\text{enc}}^\alpha$, as explained above. Integrating over the remaining loop time t_{L+1} (instead of the dwell time) and summing over all $\vec{v}$ we obtain the complete expansion of the conductance

$$\begin{aligned} &\text{Tr}(t^\dagger t) = \frac{N_1 N_2}{N} \\ &= \frac{N_1 N_2}{T_H} \sum_{\vec{v}} \mathcal{N}(\vec{v}) \left(\prod_{i=1}^{L+1} \int_0^\infty dt_i e^{-\frac{N}{T_H} t_i} \right) \\ &\times \left\langle \int \frac{d^{L-V} u d^{L-V} s}{\Omega^{L-V}} \prod_{\alpha=1}^V \frac{e^{-\frac{N}{T_H} t_{\text{enc}}^\alpha + \frac{1}{\hbar} \sum_j s_{\alpha j} u_{\alpha j}}}{t_{\text{enc}}^\alpha} \right\rangle \\ &= \frac{N_1 N_2}{N} \sum_{n=1}^\infty \frac{1}{N^n} \sum_{\vec{v}}^{(L-V=n)} (-1)^V \mathcal{N}(\vec{v}). \end{aligned} \quad (7)$$

Each loop is represented by one loop-time integral as a factor and each encounter by an $(l-1)$ fold (s, u) -integral. The (s, u) -integrals are done as above, $e^{-\frac{N}{T_H} t_{\text{enc}}^\alpha} \rightarrow -\frac{N}{T_H} t_{\text{enc}}^\alpha$ and $(\frac{1}{\Omega} \int ds du e^{isu/\hbar})^{L-V} \rightarrow T_H^{V-L}$.

In the last member of (7) the sum over the possible vectors $\vec{v}$ has been split into two, one with $\vec{v}$ restricted as $L - V = n$ and a subsequent one over the integer n . The partial sum with fixed n can be carried out with the help of a recursion relation for the $\mathcal{N}(\vec{v})$ obtained in [3] and reads (see the first technical note below)

$$\sum_{\vec{v}}^{(L-V=n)} (-1)^V \mathcal{N}(\vec{v}) = (1 - 2/\beta)^n. \quad (8)$$

Upon inserting that partial sum into (7) we recover the full $1/N$ -expansion (1) of the conductance, now for a single device rather from an ensemble average.

The *transition between the orthogonal and unitary symmetry class in a weak magnetic field* has been discussed in Ref. [4, 7, 12]. Saito and Nagao [13] have recently accounted for a magnetic field in the semiclassical τ -expansion of the spectral form factor, up to τ^3 . We here extend their ideas to the $1/N$ -expansion of

the conductance. The additional action accumulated in the magnetic field $\mathbf{B}$ on some trajectory interval CD is $\theta_{CD} = \frac{e}{c} \int_C^D \mathbf{A} \mathbf{v} dt$ with $\mathbf{A}$ the vector potential. If the dynamics is chaotic the effect of the field has been shown to be representable by Gaussian white noise [4] which yields the average $\langle e^{i2\theta_{CD}/\hbar} \rangle_G = e^{-b t_{CD}}$; here b , scaling like θ_{CD}^2 , is $(B/\hbar)^2$ times a system specific parameter. We shall see presently that net contributions of that type come both from loops and encounter stretches.

We consider a pair (γ, γ') and denote by θ_i the magnetic action gained on the i -th loop with duration t_i in γ . If that loop is passed in the same direction by γ and γ' that action cancels in $(S_\gamma - S_{\gamma'})/\hbar$; otherwise it is doubled. Defining a quantity δ_i equal to 0 and 1 in the first and second case, respectively, we can write as $\langle e^{i2\delta_i \theta_i/\hbar} \rangle_G = e^{-b\delta_i t_i}$ the additional factor in the contribution of the pair to the conductance associated with the loop “ i ”. Now consider an encounter “ α ” and denote by θ_α the magnetic action accumulated on a single stretch of this encounter passed in some (“positive”) direction. The magnetic action gained on all stretches of “ α ” in γ will be $\sigma_\alpha \theta_\alpha$ with σ_α the difference between the number of the encounter stretches traversed in the positive and negative direction. The corresponding action for the partner γ' will be $\sigma'_\alpha \theta_\alpha$; generally $\sigma'_\alpha \neq \sigma_\alpha$. Averaging yields the factor $\langle e^{i2\mu_\alpha \theta_\alpha/\hbar} \rangle_G = e^{-\mu_\alpha^2 b t_{\text{enc}}^\alpha}$ in the contribution of the orbit pair, with the integer $\mu_\alpha = (\sigma_\alpha - \sigma'_\alpha)/2$.

We shall parametrize the magnetic field by $\xi = bT_H/N \propto (B/\hbar)^2$ and denote by $\nu = \sum_{i=1}^{L+1} \delta_i$ the number of loops whose sense of traversal is changed in γ' relative to γ . Finally, we refine our definition of the number of structures by introducing the numbers $\tilde{\mathcal{N}}(\vec{v}, \nu, \{\mu_\alpha\})$ of topologically different trajectory pairs with fixed $\vec{v}, \nu, \{\mu_\alpha\}$. The above result (7) is then reformulated for the magnetoconductance,

$$\begin{aligned} & \frac{N}{N_1 N_2} \left(\text{Tr}(t^\dagger t) - \frac{N_1 N_2}{N} \right) \\ &= \frac{1}{T_H} \sum_{\vec{v}, \nu, \{\mu_\alpha\}} \tilde{\mathcal{N}}(\vec{v}, \nu, \{\mu_\alpha\}) \prod_{i=1}^{L+1} \int_0^\infty dt_i e^{-(1+\xi\delta_i)t_i \frac{N}{T_H}} \\ & \times \left\langle \frac{d^{L-V} u d^{L-V} s}{\Omega^{L-V}} \prod_{\alpha=1}^V \frac{e^{-\frac{N}{T_H}(1+\xi\mu_\alpha^2)t_{\text{enc}}^\alpha + \frac{1}{\hbar} \sum_j s_{\alpha j} u_{\alpha j}}}{t_{\text{enc}}^\alpha} \right\rangle \\ &= \sum_{n=1}^\infty \frac{1}{N^n} \sum_{\vec{v}, \nu, \{\mu_\alpha\}} \frac{(-1)^V \tilde{\mathcal{N}}(\vec{v}, \nu, \{\mu_\alpha\}) \prod_{\alpha=1}^V (1 + \xi\mu_\alpha^2)}{(1 + \xi)^\nu}. \end{aligned} \quad (9)$$

Obviously, the limiting case $\xi \rightarrow 0$ yields the orthogonal-class conductance (7) due to $\sum_{\nu, \{\mu_\alpha\}}^{\vec{v} \text{ fixed}} \tilde{\mathcal{N}}(\vec{v}, \nu, \{\mu_\alpha\}) = \mathcal{N}(\vec{v})$; the opposite limit $\xi \rightarrow \infty$ is less transparent. The coefficients of the resulting $1/N$ -expansion for low orders n are easily obtained by checking the contributing structures of orbit pairs for their values of $\nu, \{\mu_\alpha\}$ and counting the multiplicities $\tilde{\mathcal{N}}(\vec{v}, \nu, \{\mu_\alpha\})$; we get

$$\text{Tr}(t^\dagger t) = \frac{N_1 N_2}{N} \left[1 - \frac{1}{N} \frac{1}{(1 + \xi)} + \frac{1}{N^2} \frac{1}{(1 + \xi)^2} \right] \quad (10)$$

$$- \frac{1}{N^3} \frac{\xi^4 + 4\xi^3 + 13\xi^2 + 2\xi + 1}{(1 + \xi)^5} + \dots \Big].$$

It is seen that when $\xi \rightarrow \infty$ we do recover the unitary-class conductance. Order by order, our expression (10) coincides with the RMT-based expansion of the integral obtained in [9]. For $n > 3$, it appears that our semiclassical sum (9) is easier to evaluate than the corresponding term in [9]. The expansion coefficients up to order $n = 8$ are listed in the second technical note below.

Technical note I: The pairs of open trajectories of interest here and pairs of periodic orbits with the same $\vec{v}$ have the same number $\mathcal{N}(\vec{v})$ of structures. This is a matter of definition: we had associated in [3] exactly as many structures with a single periodic orbit pair, as there are non-equivalent trajectory pairs produced by cutting one of the orbit loops. We may thus use the recurrence relations for the number of periodic orbit structures $\mathcal{N}(\vec{v})$ to find the sum (8). We need the special case incorporating equations (44,60,30) of the online version of [3],

$$\begin{aligned} \frac{2v_2}{L(\vec{v})} \mathcal{N}(\vec{v}) &- \frac{1}{L(\vec{v}) - 1} \sum_{k \geq 2} (k+1) v'_{k+1} \mathcal{N}(\vec{v}') \\ &= - \left(1 - \frac{2}{\beta} \right) \mathcal{N}(\vec{v}''). \end{aligned} \quad (11)$$

Here $\vec{v}'$ (denoted $\vec{v}^{[k, 2 \rightarrow k+1]}$ in [3]) is the vector obtained from $\vec{v}$ by decreasing the components v_2 and v_k by 1 and increasing v_{k+1} by 1; the vector $\vec{v}''$ ($\vec{v}^{[2 \rightarrow]}$ of [3]) is obtained by decreasing v_2 by 1. We multiply both parts of (11) by $(-1)^{V(\vec{v})}$ and sum over all $\vec{v}$ with fixed difference $L(\vec{v}) - V(\vec{v}) = n$. In the double sum over $\vec{v}, k$ we replace $\vec{v}$ by $\vec{v}'$ as the summation variable and use $L(\vec{v}') = L(\vec{v}) - 1, V(\vec{v}') = V(\vec{v}) - 1$; we then drop the prime in $\vec{v}'$. Similarly, we switch in the right hand side to $\vec{v}''$ as the summation variable, taking into account $L(\vec{v}'') = L(\vec{v}) - 2, V(\vec{v}'') = V(\vec{v}) - 1, L(\vec{v}'') - V(\vec{v}'') = n - 1$. Dropping the double prime we obtain

$$\begin{aligned} & \sum_{\vec{v}}^{L(\vec{v}) - V(\vec{v}) = n} \frac{(-1)^{V(\vec{v})}}{L(\vec{v})} \mathcal{N}(\vec{v}) \left[2v_2 + \sum_{k \geq 2} (k+1) v_{k+1} \right] \\ &= \left(1 - \frac{2}{\beta} \right) \sum_{\vec{v}}^{L(\vec{v}) - V(\vec{v}) = n-1} (-1)^{V(\vec{v})} \mathcal{N}(\vec{v}). \end{aligned} \quad (12)$$

The expression in the square brackets is by definition $L(\vec{v})$ and thus cancels with the denominator. The resulting recursion relation $S_n = (1 - 2/\beta) S_{n-1}$ for the sum $S_n = \sum_{\vec{v}}^{L-V=n} (-1)^V \mathcal{N}(\vec{v})$ is solved by $S_n = S_0 (1 - 2/\beta)^n$.

We conclude $S_n = 0$ for all $n \geq 1$ in the unitary case. For the orthogonal case we have to fix the constant factor S_0 . Now for $n = 1$ we have but one 2-encounter ($v_2 = 1$, all other components of $\vec{v}$ zero) and a single structure, i.e. $\mathcal{N}(\vec{v}) = 1$ such that $S_1 = -1$ and thus $S_0 = 1$.

Technical note II: We here extend the $1/N$ expansion (10) of the conductance in the presence of a magnetic field. Defining the expansion coefficients as

$$\text{Tr}(t^\dagger t) = \frac{N_1 N_2}{N} \sum_{n=0}^{\infty} \frac{C_n(\xi)}{N^n}, \quad (13)$$

and introducing the abbreviations $q_k(\xi) \equiv 1 + k^2 \xi$, $p(\xi) \equiv 1/(1 + \xi) = q_1^{-1}$ we have, by numerically determining the numbers $\tilde{\mathcal{N}}(\bar{\nu}, \nu, \{\mu_\alpha\})$,

$$\begin{aligned} C_0 &= 1, \quad C_1 = -p, \quad C_2 = p^2, \\ C_3 &= -p + q_3 p^4 - q_2^2 p^5, \\ C_4 &= p^2 - p^4 - 4q_3 p^5 + 5q_2^2 p^6, \\ C_5 &= -p + q_3 p^4 - q_2^2 p^5 + 8q_5 p^6 - 24q_2 q_4 p^7 \\ &\quad - 12q_3^2 p^7 + 49q_2^2 q_3 p^8 - 21q_2^4 p^9, \\ C_6 &= p^2 - p^4 - 4q_3 p^5 + 5q_2^2 p^6 + 4p^6 + 16q_3 p^7 \\ &\quad - 48q_5 p^7 + 168q_2 q_4 p^8 - 21q_2^2 p^8 + 84q_3^2 p^8 \\ &\quad - 392q_2^2 q_3 p^9 + 189q_2^4 p^{10}, \\ C_7 &= -p + p^4 q_3 - p^5 q_2^2 + 8p^6 q_5 - 12p^7 q_3^2 - 24p^7 q_2 q_4 \\ &\quad + 180p^8 q_7 + 49p^8 q_2^2 q_3 - 21p^9 q_2^4 - 608p^9 q_3 q_5 \end{aligned}$$

$$\begin{aligned} &-276p^9 q_4^2 - 720p^9 q_2 q_6 + 1728p^{10} q_2^2 q_5 \\ &+ 2952p^{10} q_2 q_3 q_4 + 464p^{10} q_3^3 - 3240p^{11} q_2^3 q_4 \\ &- 4440p^{11} q_2^2 q_3^2 + 5445p^{12} q_2^4 q_3 - 1485p^{13} q_2^6, \\ C_8 &= p^2 + 4p^6 - p^4 - 36p^8 + 216p^{10} q_2^2 - 21p^8 q_2^2 \\ &- 1485p^{12} q_2^4 + 5p^6 q_2^2 + 189p^{10} q_2^4 \\ &+ 19305p^{14} q_2^6 - 160p^9 q_3 + 16p^7 q_3 \\ &+ 2880p^{11} q_2^2 q_3 - 4p^5 q_3 - 392p^9 q_2^2 q_3 \\ &- 65340p^{13} q_2^4 q_3 - 1080p^{10} q_2 q_4 + 168p^8 q_2 q_4 \\ &+ 35640p^{12} q_2^3 q_4 - 624p^{10} q_3^2 + 84p^8 q_3^2 \\ &+ 48840p^{12} q_2^2 q_3^2 + 288p^9 q_5 - 48p^7 q_5 \\ &- 17280p^{11} q_2^2 q_5 - 29520p^{11} q_2 q_3 q_4 - 4640p^{11} q_3^3 \\ &+ 6480p^{10} q_2 q_6 + 2484p^{10} q_4^2 + 5472p^{10} q_3 q_5 \\ &- 1440p^9 q_7. \end{aligned}$$

A recursive treatment of all orders in the expansion (13) is called for.

We are indebted to Dmitry Savin for useful discussions and the Sonderforschungsbereich SFB/TR12 of the Deutsche Forschungsgemeinschaft for financial support.

-
- [1] R. A. Jalabert, in *Proceedings of the International School of Physics "Enrico Fermi" Course CXLIII*, edited by G. Casati, I. Guarneri, and U. Smilansky, p. 145 (IOS Press, Amsterdam, 2000)
 - [2] K. Richter, *Semiclassical Theory of Mesoscopic Quantum Systems* (Springer, 2000)
 - [3] S. Müller, S. Heusler, P. Braun, F. Haake and A. Altland, Phys. Rev. Lett. **93** (2004) 014103-1 and nlin.CD/0503052 and references therein
 - [4] K. Richter, M. Sieber, Phys. Rev. Lett. **89**, 206801-1 (2002)
 - [5] R. Landauer, J. Res. Dev. **1**, 223 (1957), and D. S. Fisher, P. A. Lee, Phys. Rev. B **23**, R6851 (1981)
 - [6] C. W. J. Beenakker, H. van Houten, *Solid State Physics* **44**, 1 (Academic Press, New York, 1991)
 - [7] C. W. J. Beenakker, Rev. Mod. Phys. **69**, 731 (1997)
 - [8] R. A. Jalabert, J. L. Pichard, and C. W. J. Beenakker, Europhys. Lett. **27**, 255 (1994); H. U. Baranger and P. A. Mello, Phys. Rev. Lett. **73**, 142 (1994)
 - [9] Z. Pluhar, H.A. Weidenmüller, J.A. Zuk, C.H. Lewenkopf, F.J. Wegner, Ann. of Phys. **243**, 1 64(1995)
 - [10] J. H. Hannay, A. M. Ozorio de Almeida, J. Phys. **A17**, 3429 (1984)
 - [11] In Ref. [4] the self-crossings of a trajectory were allowed to approach the cavity openings which means in our language that half of each encounter stretch could stick out into the leads; this is equivalent to replacing $(T - 2t_{\text{enc}})^2$ by $(T - t_{\text{enc}})^2$ in (4); in addition, the increase of the survival probability due to the self-encounter was not taken into account. The two differences to our procedure cancel in their effect on the lowest off-diagonal term $G^{(1)}$.
 - [12] H. U. Baranger, R. A. Jalabert, and A. D. Stone, Phys. Rev. Lett. **70**, 3876 (1993)
 - [13] K. Saito, T. Nagao, nlin.CD/0505055